

PACKING 3-VERTEX PATHS IN CLAW-FREE GRAPHS

Alexander Kelmans

University of Puerto Rico, San Juan, Puerto Rico
Rutgers University, New Brunswick, New Jersey

Abstract

A Λ -factor of a graph G is a spanning subgraph of G whose every component is a 3-vertex path. Let $v(G)$ be the number of vertices of G . A graph is *claw-free* if it does not have a subgraph isomorphic to $K_{1,3}$. Our results include the following. Let G be a 3-connected claw-free graph, $x \in V(G)$, $e = xy \in E(G)$, and L a 3-vertex path in G . Then (c1) if $v(G) \equiv 0 \pmod 3$, then G has a Λ -factor containing (avoiding) e , (c2) if $v(G) \equiv 1 \pmod 3$, then $G - x$ has a Λ -factor, (c3) if $v(G) \equiv 2 \pmod 3$, then $G - \{x, y\}$ has a Λ -factor, (c4) if $v(G) \equiv 0 \pmod 3$ and G is either cubic or 4-connected, then $G - L$ has a Λ -factor, and (c5) if G is cubic and E is a set of three edges in G , then $G - E$ has a Λ -factor if and only if the subgraph induced by E in G is not a claw and not a triangle.

Keywords: claw-free graph, cubic graph, Λ -packing, Λ -factor.

1 Introduction

We consider undirected graphs with no loops and no parallel edges. All notions and facts on graphs, that are used but not described here, can be found in [1, 2, 12].

Given graphs G and H , an H -packing of G is a subgraph of G whose every component is isomorphic to H . An H -packing P of G is called an H -factor if $V(P) = V(G)$. The H -packing problem, i.e. the problem of finding in G an H -packing, having the maximum number of vertices, turns out to be NP -hard if H is a connected graph with at least three vertices [3]. Let Λ denote a 3-vertex path. In particular, the Λ -packing problem is NP -hard. Moreover, this problem remains NP -hard even for cubic graphs [6].

Although the Λ -packing problem is NP -hard, i.e. possibly intractable in general, this problem turns out to be tractable for some natural classes of graphs (see, for example, **1.10** below). It would be also interesting to find polynomial-time algorithms that would provide a good approximation solution for the problem (e.g. **1.1** and **1.11** below). In each case the corresponding packing problem is polynomially solvable.

Let $v(G)$ and $\lambda(G)$ denote the number of vertices and the maximum number of disjoint 3-vertex paths in G , respectively. Obviously $\lambda(G) \leq \lfloor v(G)/3 \rfloor$.

In [5, 11] we answered the following natural question:

How many disjoint 3-vertex paths must a cubic n -vertex graph have?

1.1 *If G is a cubic graph, then $\lambda(G) \geq \lceil v(G)/4 \rceil$. Moreover, there is a polynomial time algorithm for finding a Λ -packing having at least $\lceil v(G)/4 \rceil$ components.*

Obviously if every component of G is K_4 , then $\lambda(G) = v(G)/4$. Therefore the bound in **1.1** is sharp.

Let \mathcal{G}_2^3 denote the set of graphs with each vertex of degree 2 or 3. In [5] we answered (in particular) the following question:

How many disjoint 3-vertex paths must an n -vertex graph from \mathcal{G}_2^3 have?

1.2 *Suppose that $G \in \mathcal{G}_2^3$ and G has no 5-vertex components. Then $\lambda(G) \geq v(G)/4$.*

Obviously **1.1** follows from **1.2** because if G is a cubic graph, then $G \in \mathcal{G}_2^3$ and G has no 5-vertex components.

In [5] we also gave a construction that allowed to prove the following:

1.3 *There are infinitely many connected graphs for which the bound in **1.2** is attained. Moreover, there are infinitely many subdivisions of cubic 3-connected graphs for which the bound in **1.2** is attained.*

The next interesting question is:

How many disjoint 3-vertex paths must a cubic connected graph have?

In [7] we proved the following. Let \mathcal{C}_n denote the set of connected cubic graphs with n vertices.

1.4 *Let $\lambda_n = \min\{\lambda(G)/v(G) : G \in \mathcal{C}_n\}$. Then for some $c > 0$,*

$$\frac{3}{11}(1 - \frac{c}{n}) \leq \lambda_n \leq \frac{3}{11}(1 - \frac{1}{n^2}).$$

The similar question for cubic 2-connected graphs is still open:

1.5 Problem. *How many disjoint 3-vertex paths must a cubic 2-connected graph have?*

It is known that

1.6 *There are infinitely many 2-connected and cubic graphs G such that $\lambda(G) < \lfloor v(G)/3 \rfloor$.*

Some such graph sequences were constructed in [10] to provide 2-connected counterexamples to Reed's domination conjecture. Reed's conjecture claims that if G is a connected cubic graph, then $\gamma(G) \leq \lceil v(G)/3 \rceil$, where $\gamma(G)$ is the dominating number of G (i.e. the size of a minimum vertex subset X in G such that every vertex in $G - X$ is adjacent to a vertex in X). In particular, a graph sequence $(R_k : k \geq 3)$ in [10] is such that each R_k is a cubic graph of connectivity two and $\gamma(G)/v(G) = \frac{1}{3} + \frac{1}{60}$. Obviously, $\gamma(G) \leq v(G) - 2\lambda(G)$. Therefore $\lambda(R_k)/v(R_k) \leq \frac{13}{40}$.

The questions arise whether the claim of **1.6** is true for cubic 2-connected graphs having some additional properties. For example,

1.7 Problem. *Is $\lambda(G) = \lfloor v(G)/3 \rfloor$ true for every 2-connected, cubic, bipartite, and planar graph ?*

In [8] we answered the question in **1.7** by giving a construction that provides infinitely many 2-connected, cubic, bipartite, and planar graphs such that $\lambda(G) < \lfloor v(G)/3 \rfloor$.

As to cubic 3-connected graphs, an old open question here is:

1.8 Problem. *Is the following claim true ?
If G is a 3-connected and cubic graph, then $\lambda(G) = \lfloor v(G)/3 \rfloor$.*

In [9] we discuss Problem **1.8** and show, in particular, that the claim in **1.8** is equivalent to some seemingly much stronger claims. Here are some results of this kind.

1.9 [9] *The following are equivalent for cubic 3-connected graphs G :*

- (z1) $v(G) \equiv 0 \pmod 6 \Rightarrow G$ has a Λ -factor,
- (z2) $v(G) \equiv 0 \pmod 6 \Rightarrow$ for every $e \in E(G)$ there is a Λ -factor of G avoiding e ,
- (z3) $v(G) \equiv 0 \pmod 6 \Rightarrow$ for every $e \in E(G)$ there is a Λ -factor of G containing e ,
- (z4) $v(G) \equiv 0 \pmod 6 \Rightarrow G - X$ has a Λ -factor for every $X \subseteq E(G)$, $|X| = 2$,
- (z5) $v(G) \equiv 0 \pmod 6 \Rightarrow G - L$ has a Λ -factor for every 3-vertex path L in G ,
- (t1) $v(G) \equiv 2 \pmod 6 \Rightarrow G - \{x, y\}$ has a Λ -factor for every $xy \in E(G)$,
- (f1) $v(G) \equiv 4 \pmod 6 \Rightarrow G - x$ has a Λ -factor for every $x \in V(G)$,
- (f2) $v(G) \equiv 4 \pmod 6 \Rightarrow G - \{x, e\}$ has a Λ -factor for every $x \in V(G)$ and $e \in E(G)$.

From **1.9** it follows that if the claim in Problem **1.8** is true, then Reed's domination conjecture is true for 3-connected cubic graphs.

There are some interesting results on the Λ -packing problem for so called claw-free graphs. A graph is called *claw-free* if it contains no induced subgraph isomorphic to $K_{1,3}$ (which is called a *claw*).

A vertex x of a block B in G is called a *boundary vertex of B* if x belongs to another block of G . If B has exactly one boundary vertex, then B is called an *end-block* of G . Let $eb(G)$ denote the number of end-blocks of G .

1.10 [4] *Suppose that G is a claw-free graph and G is either 2-connected or connected with exactly two end-blocks. Then $\lambda(G) = \lfloor v(G)/3 \rfloor$.*

1.11 [4] *Suppose that G is a connected claw-free graph and $eb(G) \geq 2$. Then $\lambda(G) \geq \lfloor (v(G) - eb(G) + 2)/3 \rfloor$, and this lower bound is sharp.*

Obviously the claim in **1.10** about connected claw-free graphs with exactly two end-blocks follows from **1.11**.

In this paper (see Section 2) we give some more results on the Λ -packings in claw-free graphs. We show, in particular, the following:

- (c1) all claims in **1.9** except for (z5) are true for 3-connected claw-free graphs and (z5) is true for cubic, 2-connected, and claw-free graphs distinct from K_4 (see **2.15** below),
- (c2) if G is a 3-connected claw-free graph and $v(G) \equiv 0 \pmod 3$, then for every edge e in G there exists a Λ -factor of G containing e (see **2.7**),
- (c3) if G is a 2-connected claw-free graph and $v(G) \equiv 0 \pmod 3$, then for every edge e in G there exists a Λ -factor of G avoiding e , i.e. $G - e$ has a Λ -factor (see **2.8**),
- (c4) if G is a 2-connected claw-free graph and $v(G) \equiv 1 \pmod 3$, then $G - x$ has a Λ -factor for every vertex x in G (see **2.13**),
- (c5) if G is a 3-connected claw-free graph and $v(G) \equiv 2 \pmod 3$, then $G - \{x, y\}$ has a Λ -factor for every edge xy in G (see **2.3**),
- (c6) if G is a 3-connected, claw-free, and cubic graph with $v(G) \geq 6$ or a 4-connected claw-free graph, then for every 3-vertex path L in G there exists a Λ -factor containing L , i.e. $G - L$ has a Λ -factor (see **2.1** and **2.6**),
- (c7) if G is a cubic, 3-connected, and claw-free graph with $v(G) \geq 6$ and E is a set of three edges in G , then $G - E$ has a Λ -factor if and only if the subgraph induced by E in G is not a claw and not a triangle (see **2.10**),
- (c8) if G is a 3-connected claw-free graph, $v(G) \equiv 1 \pmod 3$, $x \in V(G)$, and $e \in E(G)$, then $G - \{x, e\}$ has a Λ -factor (see **2.14**).

2 Main results

Theorems **1.10** and **1.11** above describe some properties of maximum Λ -packings in claw-free graphs. In this section we establish some more properties of Λ -packings in claw-free graphs.

Let G be a graph and B be a block of G , and so B is either 2-connected or consists of two vertices and one edge. As above, a vertex x of B is called a *boundary vertex* of B if x belongs to another block of G , and an *inner vertex* of B , otherwise. If B has exactly one boundary vertex, then B is called an *end-block* of G .

Let F be a graph, $x \in V(F)$, and $X = \{x_1, x_2, x_3\}$ be the set of vertices in F adjacent to x . Let T be a triangle, $V(T) = \{t_1, t_2, t_3\}$, and $V(F) \cap V(T) = \emptyset$. Let $G = (F - x) \cup T \cup \{x_i t_i : i \in \{1, 2, 3\}\}$. We say that G is obtained from F by replacing a vertex x by a triangle. Let F^Δ denote the graph obtained from a cubic graph F by replacing each vertex of F by a triangle. Obviously, F^Δ is claw-free, every vertex belongs to exactly one triangle, and every edge belongs to at most one triangle in F^Δ .

2.1 Let G' be a cubic 2-connected graph and G be the graph obtained from G' by replacing each vertex v of G' by a triangle Δ_v . Let L be a 3-vertex path in G . Then

(a) $G - L$ has a Λ -factor.

Moreover,

(a1) if L induces a triangle in G , then G has a Λ -factor R containing L and such that each component of R induces a triangle

(a2) if L does not induce a triangle in G , then G has a Λ -factor R containing L and such that no component of R induces a triangle, and

(a3) if L does not induce a triangle in G , then G has a Λ -factor, containing L and a component that induces a triangle.

Proof Let $L = xzz_1$. Let E' be the set of edges in G that belong to no triangle. Obviously, there is a natural bijection $\alpha : E(G') \rightarrow E'$. Since each vertex of G belongs to exactly one triangle, we can assume that xz belongs to a triangle $T = xzs$.

(p1) Suppose that L induces a triangle in G , and so $s = z_1$. Obviously the union of all triangles in G contains a Λ -factor, say P , of G and $L \subset P$. Therefore claim (a1) is true.

(p2) Now suppose that L does not induce a triangle in G , and so $s \neq z_1$. Let $\bar{s} = ss_1$ and $\bar{z} = zz_1$ be the edges of G not belonging to T , and therefore belonging to no triangles in G . Hence $\bar{s} = \alpha(\bar{s}')$ and $\bar{z} = \alpha(\bar{z}')$, where $\bar{s}' = s's'_1$ and $\bar{z}' = z'z'_1$ are edges in G' , and $s' = z'$. Since every vertex in G belongs to exactly one triangle, clearly $s_1 \neq z_1$.

(p2.1) We prove (a2). By using Tutte's criterion for a graph to have a perfect matching, it is easy to prove the following:

CLAIM. If A is a cubic 2-connected graph, then for every 3-vertex path J of A there exists a 2-factor of A containing J .

By the above CLAIM, G' has a 2-factor F' containing 3-vertex path $S' = s'_1s'z'_1$. Let C' be the (cycle) component of F' containing S' . If Q' is a (cycle) component of F' , then let Q be the subgraph of G , induced by the edge subset $\{\alpha(e) : e \in E(Q')\} \cup \{E(\Delta_v) : v \in V(Q')\}$. Obviously $v(Q) = 0 \pmod 3$ and Q has a (unique) Hamiltonian cycle $H(Q)$. Also the union F of all Q 's is a spanning subgraph of G and each Q is a component of F . Moreover, if C is the component in F , corresponding to C' , then $L \subset H(C)$. Therefore each $H(Q)$ has a Λ -factor $P(Q)$, such that no component of $P(Q)$ induces a triangle, and $H(C)$ has a (unique) Λ -factor $P(C)$, such that $L \subset P(C)$ and no component of $P(C)$ induces a triangle. The union of all these Λ -factors is a Λ -factor P of G containing L and such that no component of P induces a triangle. Therefore (a2) holds.

(p2.2) Now we prove (a3). Since G' is 2-connected and cubic, there is a cycle C' in G' such that $V(C') \neq V(G')$ and C' contains $S' = s'_1s'z'_1$. Let, as above, C be the subgraph of G , induced by the edge subset $\{\alpha(e) : e \in E(C')\} \cup \{E(\Delta_v) : v \in V(C')\}$. Obviously, $v(C) = 0 \pmod 3$, C has a (unique) Hamiltonian cycle H , and $L \subset H$. Therefore H has a (unique) Λ -factor $P(C)$ containing L . Since $V(C') \neq V(G')$, we have $V(G' - C') \neq \emptyset$.

Therefore $G - C$ has a triangle. Moreover, every vertex v in $G - C$ belongs to a unique triangle Δ_v , and therefore as in (p1), $G - C$ has a Λ -factor Q whose every component induces a triangle in $G - C$. Then $P(C) \cup Q$ is a required a Λ -factor in G . \square

Theorem 2.1 is not true for a cubic, 2-connected, and claw-free graph F with an edge xy belonging to two triangles T_i with $V(T_i) = \{x, y, z_i\}$ because $L = z_1 x z_2$ is a 3-vertex path in F and y is an isolated vertex in $F - L$.

2.2 Suppose that G is a 2-connected claw-free graph, $v(G) = 2 \pmod 3$, and $x \in V(G)$. Then there exist at least two edges xz_1 and xz_2 in G such that each $G - xz_i$ is connected and has a Λ -factor.

Proof (uses 1.10). We need the following simple facts.

CLAIM 1. Let G be a 2-connected graph and, $x \in V(G)$. Then there exist at least two edges xs_1 and xs_2 in G such that each $G - xs_i$ is connected.

CLAIM 2. Let G be a claw-free graph, B is a 2-connected block of G , and x is a boundary vertex of B . Then $B - x$ is either 2-connected or has exactly one edge.

By CLAIM 1, G has an edge xy such that $G - \{x, y\}$ is connected. If $G - \{x, s\}$ for every $xs \in E(G)$, then by CLAIM 1, we are done. Therefore we assume that $G - \{x, y\}$ is connected but has no Λ -factor.

Then by 1.10, $G - \{x, y\}$ has at least three end-blocks, say B_i , $i \in \{1, \dots, k\}$, $k \geq 3$. Let b'_i be the boundary vertex of B_i . Let V_i be the set of vertices in $\{x, y\}$ adjacent to the interior of B_i and \mathcal{B}_v be the set of the end-blocks in $G - \{x, y\}$ whose interior is adjacent to $v \in \{x, y\}$. Since G is 2-connected, each $|V_i| \geq 1$. Since G is claw-free, each $|\mathcal{B}_v| \leq 2$. Since $k \geq 3$, $|\mathcal{B}_v| = 2$ for some $v \in \{x, y\}$, say for $v = x$ and $\mathcal{B}_x = \{B_1, B_2\}$. Let $xb_i \in E(G)$, where b_i is an interior vertex of B_i , $i \in \{1, 2\}$, and let $xb_j \in E(G)$, where b_j is an interior vertex of B_j , $j \geq 3$. Since G is claw-free, $\{x, y, b_1, b_2\}$ does not induce a claw in G . Therefore $y b_2 \in E(G)$. If $k \geq 4$, then $\{y, b_2, b_3, b_4\}$ induces a claw in G , a contradiction. Thus $k = 3$ and $\mathcal{B}_y = \{B_2, B_3\}$.

Suppose that $v(B_s) = 0 \pmod 3$ for some $s \in \{1, 2, 3\}$. Then B_s is 2-connected. Since B_s is claw-free, by 1.10, B_s has a Λ -factor, say P . Since $v(G) = 2 \pmod 3$, we have $v(G - \{x, y, B_s\}) = 0 \pmod 3$. By CLAIM 2, $G - \{x, y, B_s\}$ is claw-free, connected and has at most two end-blocks. Then by 1.10, $G - \{x, y, B_s\}$ has Λ -factor, say Q . Therefore $P \cup Q$ is a Λ -factor of $G - \{x, y\}$, a contradiction.

Suppose that $v(B_r) = 1 \pmod 3$ for some $r \in \{1, 2, 3\}$. Then B_r is 2-connected. Obviously $v(B_r - b'_r) = 0 \pmod 3$ and claw-free. By CLAIM 2, $B_r - b'_r$ is 2-connected. Then by 1.10, B_r has a Λ -factor, say P . Obviously $G - \{x, y, B_r - b'_r\}$ is claw-free, connected and has at most two end-blocks. Then by 1.10, $G - \{x, y, B_s\}$ has Λ -factor, say Q . Therefore $P \cup Q$ is a Λ -factor of $G - \{x, y\}$, a contradiction.

Now suppose that $v(B_i) = 2 \pmod 3$ for every $i \in \{1, 2, 3\}$. By CLAIM 2, either $B_i - \{b_i, b'_i\}$ is 2-connected or $v(B_i - \{b_i, b'_i\}) = 0$ for every $i \in \{1, 2, 3\}$. In both cases

by the arguments similar to that above, $G - \{x, b_i\}$ has a Λ -factor for $i \in \{1, 2\}$ and $G - \{y, b_i\}$ has a Λ -factor for $i \in \{2, 3\}$. \square

From **2.2** we have for 3-connected claw-free graphs the following stronger result with a simpler proof.

2.3 *Suppose that G is a 3-connected claw-free graph, $v(G) = 2 \pmod 3$, and $xy \in E(G)$. Then $G - \{x, y\}$ has a Λ -factor.*

Proof (uses **1.10**). Let $G' = G - \{x, y\}$. Since G is 3-connected, G' is connected. By **1.10**, it suffices to prove that G' has at most two end-blocks. Suppose, on the contrary, that G' has at least three end-blocks. Let B_i , $i \in \{1, 2, 3\}$, be some three blocks of G' . Since G is 3-connected, for every block B_i and every vertex $v \in \{x, y\}$ there is an edge vb_i , where b_i is an inner vertex of B_i . Then $\{v, b_1, b_2, b_3\}$ induces a claw in G , a contradiction. \square

As we have seen in the proof of **2.2**, the claim of **2.3** is not true for claw-free graphs of connectivity two.

2.4 *Suppose that G is a 3-connected claw-free graph, $v(G) = 0 \pmod 3$, and $xy \in E(G)$. Then there exist at least two 3-vertex paths L_1 and L_2 in G centered at y , containing xy , and such that each $G - L_i$ is connected and has a Λ -factor.*

Proof (uses **1.10**). We need the following simple fact.

CLAIM 1. *Let G be a 3-connected graph, $x \in V(G)$, and $xy \in E(G)$. Then there exist two 3-vertex paths L_1 and L_2 in G centered at y , containing xy , and such that each $G - L_i$ is connected.*

By CLAIM 1, G has a 3-vertex path $L = xyz$ such that $G - L$ is connected. If every such 3-vertex path belongs to a Λ -factor of G , then by CLAIM 1, we are done. Therefore we assume that $G - L$ is connected but has no Λ -factor. Then by **1.10**, $G - L$ has at least three end-blocks, say B_i , $i \in \{1, \dots, k\}$, $k \geq 3$. Let b'_i be the boundary vertex of B_i . Let V_i be the set of vertices in L adjacent to inner vertices of B_i and \mathcal{B}_v be the set of the end-blocks in $G - L$ having an inner vertex adjacent to v in $V(L)$. Since G is 3-connected, each $|V_i| \geq 2$. Since G is claw-free, each $|\mathcal{B}_v| \leq 2$. It follows that $k = 3$, each $|V_i| = 2$, each $|\mathcal{B}_v| = 2$, as well as all V_i 's are different and all \mathcal{B}_v 's are different. Let $s^1 = z$, $s^2 = x$, $s^3 = y$, and $S = \{s^1, s^2, s^3\}$. We can assume that $V_i = S - s^i$, $i \in \{1, 2, 3\}$. Then for every vertex $s^j \in V_i$ there is a vertex b_i^j in $B_i - b'_i$ adjacent to s^j , where $\{b_i^j : s^j \in V_i\}$ has exactly one vertex if and only if $B_i - b'_i$ has exactly one vertex. Let $L_i = s^2 s^3 b_i$, where $b_i = b_i^3$.

By **1.10**, it suffices to show that each $G - L_i$ is connected and has at most two end-blocks.

Let $i = 1$. If $B_1 - b_1$ is 2-connected, then $B_1 - b_1$ and $G - L_1 - (B_1 - b'_1)$ are the two end-blocks of $G - L_1$ and we are done. If $B_1 - b_1$ is empty, then $G - L_1$ is 2-connected.

So we assume that $B_1 - b_1$ is not empty and not 2-connected. Then $B_1 - b_1$ is connected and has exactly two end-blocks, say C_1 and C_2 . Let c'_i be the boundary vertex of C_i in $B_1 - b_1$. Since G is 3-connected, each $C_i - c'_i$ has a vertex adjacent to $\{s^2, s^3\}$. We can assume that a vertex c_1 in $C_1 - c'_1$ is adjacent to s^2 . If there exists a vertex c_2 in $C_2 - c'_2$ adjacent to s^2 , then $\{s^2, b_3^2, c_1, c_2\}$ induces a claw in G , a contradiction. So suppose that no vertex in $C_2 - c'_2$ is adjacent to s^2 . Then there is a vertex c_2 in $C_2 - c'_2$ adjacent to s^3 . Then $\{s^2, s^3, b_2^3, c_2\}$ induces a claw in G , a contradiction.

Now let $i = 2$. If $B_2 - b_2$ is 2-connected, then B_1 and $G - L_2 - (B_1 - b'_1)$ are the two end-blocks of $G - L_2$ and we are done. If $B_1 - b_1$ is empty, then $G - L_2$ has two end-blocks, namely B_1 and the subgraph of G induced by $B_3 \cup s^1$. So we assume that $B_2 - b_2$ is not empty and not 2-connected. Then $B_2 - b_2$ is connected and has exactly two end-blocks, say D_1 and D_2 . Let d'_i be the boundary vertex of D_i in $B_2 - b_2$. Since G is 3-connected, each $D_i - d'_i$ has a vertex adjacent to $\{s^1, s^3\}$. We can assume that a vertex d_1 in $D_1 - d'_1$ is adjacent to s^3 . If there exists a vertex d_2 in $D_2 - d'_2$ adjacent to s^3 , then $\{s^3, d_1, d_2, b_1^3\}$ induces a claw in G , a contradiction. So suppose that no vertex in $D_2 - d'_2$ is adjacent to s^3 . Then there is a vertex d_2 in $D_2 - d'_2$ adjacent to s^1 . Then $\{s^1, s^3, b_3^1, d_2\}$ induces a claw in G , a contradiction. \square

From the proof of **2.4** we have, in particular:

2.5 *Suppose that G is a 3-connected claw-free graph. If L is a 3-vertex path and the center vertex of L has degree 3 in G , then $G - L$ is connected and has a Λ -factor in G .*

Obviously, **2.1** (a) follows from **2.5**.

From the proof of **2.4** we also have:

2.6 *Suppose that G is a 4-connected claw-free graph. Then $G - L$ is connected and has a Λ -factor for every 3-vertex path L in G .*

The claim of **2.6** may not be true for a claw-free graph of connectivity 3 if they are not cubic. A graph obtained from a claw by replacing its vertex of degree 3 by a triangle is called a *net*. Let N be a net with the three leaves v_1, v_2 , and v_3 , T a triangle with $V(T) = \{t_1, t_2, t_3\}$, and let N and T be disjoint. Let $H = N \cup T \cup \{v_i t_j : i, j \in \{1, 2, 3\}, i \neq j\}$. Then H is a 3-connected claw-free graph, $v(H) = 9$, each $d(t_i, H) = 4$, $d(x, H) = 3$ for every $x \in V(H - T)$, and $H - T = N$ has no Λ -factor. If L is a 3-vertex path in T , then $H - L = H - T$, and so $H - L$ has no Λ -factor. There are infinitely many pairs (G, L) such that G is a 3-connected, claw-free, and non-cubic graph, $v(G) \equiv 0 \pmod 3$, L is a 3-vertex path in G , and $G - L$ has no Λ -factor. By **2.9**, such a pair can be obtained from the above pair (H, L) by replacing N by any graph A with three leaves from the class \mathcal{A} (defined below before **2.9**) provided $v(A) \equiv 0 \pmod 3$.

From **2.4** we have, in particular:

2.7 Suppose that G is a 3-connected claw-free graph and $e \in E(G)$. Then

(a1) there exists a Λ -factor in G containing e and

(a2) there exists a Λ -factor in G avoiding e , i.e. $G - e$ has a Λ -factor.

The following examples show that condition “ G is a 3-connected graph” in **2.7** is essential for claim (a1). Let R be the graph obtained from two disjoint cycles A and B by adding a new vertex z , and the set of new edges $\{a_i z, b_i z : i \in \{1, 2\}\}$, where $a = a_1 a_2 \in E(A)$ and $b = b_1 b_2 \in E(B)$. It is easy to see that R is a claw-free graph of connectivity one. Furthermore, if $v(A) = 1 \pmod 3$ and $v(B) = 1 \pmod 3$, then $v(R) = 0 \pmod 3$ and R has no Λ -factor containing edge $e \in \{a, b\}$. Similarly, let Q be the graph obtained from two disjoint cycles A and B by adding two new vertices z_1 and z_2 , a new edge $e = z_1 z_2$, and the set of new edges $\{a_i z_j, b_i z_j : i, j \in \{1, 2\}\}$, where $a_1 a_2 \in E(A)$ and $b_1 b_2 \in E(B)$. It is easy to see that Q is a claw-free graph of connectivity two. Furthermore, if $v(A) = 2 \pmod 3$ and $v(B) = 2 \pmod 3$, then $v(Q) = 0 \pmod 3$ and Q has no Λ -factor containing edge e .

As to claim (a2) in **2.7**, it turns out that this claim is also true for 2-connected claw-free graphs.

2.8 Suppose that G is a 2-connected claw-free graph, $v(G) = 0 \pmod 3$, and $e \in E(G)$. Then $G - e$ has a Λ -factor.

Proof A graph H is called *minimal 2-connected* if H is 2-connected but $H - u$ is not 2-connected for every $u \in E(H)$. A *frame* of a graph G is a minimal 2-connected spanning subgraph of G . Clearly, every 2-connected graph has a frame. In [4] we describe Procedure 1 that provides an ear-assembly A of a special frame of a 2-connected claw-free graph. In particular, the last ear of A contains a Λ -packing P such that $G - P$ is also 2-connected claw-free graph. We modify Procedure 1 by replacing the first step of this procedure “Find a longest cycle G_0 in G ” by “Find a longest cycle G'_0 among all cycles C in G such that edge e either belongs to C or is a chord of C ”. Since G is 2-connected, G has a cycle containing e . Therefore a cycle G'_0 does exist. Then the resulting Procedure \mathcal{P} provides an ear-assembly of a frame of G with the property that the last ear of this frame has a Λ -packing Q such that $e \notin E(Q)$ and $G - Q$ is a 2-connected claw-free graph that may contain e .

We can use Procedure \mathcal{P} to prove our claim by induction on $v(G)$. If G is a cycle containing e , then our claim is obviously true. Procedure \mathcal{P} mentioned above guarantees the existence of a Λ -packing Q such that Q avoids e and $G - Q$ is a 2-connected claw-free graph that may contain e . Obviously $v(G - Q) = 0 \pmod 3$ and $v(G - Q) < v(G)$. By the induction hypothesis, $G - Q$ has a Λ -factor R avoiding e . Then $Q \cup R$ is a Λ -factor of G avoiding edge e . \square

We need the following fact interesting in itself. Let \mathcal{A} denote the set of graphs A with the following properties:

- (c1) A is connected,
- (c2) every vertex in A has degree at most 3,
- (c3) every vertex in A of degree 2 or 3 belongs to exactly one triangle, and
- (c4) A has exactly three vertices of degree 1 which we call the *leaves* of A .

2.9 *If $A \in \mathcal{A}$, then A has no Λ -factor.*

Proof Let $A \in \mathcal{A}$. If $v(A) \not\equiv 0 \pmod 3$, then our claim is clearly true. So we assume that $v(A) \equiv 0 \pmod 3$. We prove our claim by induction on $v(G)$. The smallest graph in \mathcal{A} is a net N with $v(N) = 6$ and our claim is obviously true for N . So let $v(A) \geq 9$. Suppose, on the contrary, that A has a Λ -factor P . Let v be a leaf of A and vx the edge incident to v . Since P is a Λ -factor in A , it has a component $L = vxy$, and so $P - L$ is a Λ -factor in $A - L$ and $d(x, A) \geq 2$. By property (c3), x belongs to a unique triangle xyz in A and $d(x, A) = 3$, and so $s \in \{y, z\}$. If $d(z, A) = 2$, then z is an isolated vertex in $A - L$, and so P is not a Λ -factor in A , a contradiction. Therefore by (c2), $d(z, A) = 3$. Therefore $A - L$ satisfies (c2), (c3), and (c4).

Suppose that $G - L$ is not connected and that the three leaves do not belong to a common component. Then $A - L$ has a component C with $v(C) \not\equiv 0 \pmod 3$, and so $A - L$ has no Λ -factor, a contradiction.

Now suppose that $A - L$ has a component C containing all three leaves of $A - L$. Then $C \in \mathcal{A}$ and $v(C) < v(A)$. By the induction hypothesis, C has no Λ -factor. Therefore $A - L$ also has no Λ -factor, a contradiction. \square

Given $E \subseteq E(G)$, let \dot{E} denote the subgraph of G induced by E .

2.10 *Suppose that G is a cubic 2-connected graph and that every vertex in G belongs to exactly one triangle (and so G is claw-free), i.e. $G = F^\Delta$, where F is a cubic 2-connected graph. Let $E \subset E(G)$ and $|E| = 3$. Then the following are equivalent:*

- (g) $G - E$ has no Λ -factor and
- (e) \dot{E} satisfies one of the following conditions:

- (e1) \dot{E} is a claw,
- (e2) \dot{E} is a triangle,
- (e3) \dot{E} has exactly two components, the 2-edge component \dot{E}_2 belongs to a triangle in G , the 1-edge component \dot{E}_1 belongs to no triangle in G , and $G - E$ is not connected, and

(e4) \dot{E} has exactly two components, in G the 2-edge component \dot{E}_2 belongs to a triangle, say T , the 1-edge component \dot{E}_1 also belongs to a triangle, say D , and \dot{E}_1, \dot{E}_2 belong to different component of $G - \{d, t\}$, where d is the edge incident to the vertex of $D - \dot{E}_1$ and t is the edge in $G - E$ incident to the isolated vertex of $T - E$.

Proof (uses 1.10, 2.1(a), and 2.9). Let $X, Y \subset E(G)$ such that X meets no triangle in G , each edge in Y belongs to a triangle in G , and no triangle in G has more than one

edge from Y , and so $X \cap Y = \emptyset$. We will use the following simple observation.

CLAIM. $G - X - Y$ has a Λ -factor P such that every component of P induces a triangle in G and if an edge y from Y is in a triangle T , then $T - y$ is a component of P .

Let $E = \{a, b, c\}$. By the above CLAIM, we can assume that edges a and b belong to the same triangle T .

(p1) We prove $(e) \Rightarrow (g)$.

Suppose that \dot{E} satisfies (e1), i.e. \dot{E} is a claw. Then $G - E$ has an isolated vertex and therefore has no Λ -factor.

Suppose that \dot{E} satisfies (e2), i.e. \dot{E} is a triangle. Then $G - E \in \mathcal{A}$. By **2.9**, $G - E$ has no Λ -factor.

Suppose that \dot{E} satisfies (e3), i.e. $G - E$ is not connected and \dot{E} has exactly two components \dot{E}_2 and \dot{E}_1 induced by $\{a, b\}$ and c , respectively, where \dot{E}_2 belongs to the triangle T but \dot{E}_1 belongs to no triangle in G . Then t is the dangling edge in $G - E$. Let S be the component in $G - E$ containing edge t . Then every vertex in S distinct from the leaf incident to t belongs to exactly one triangle. Therefore $v(S) = 1 \pmod 3$. Thus $G - E$ has no Λ -factor.

Now suppose that \dot{E} satisfies (e4). By (e4), t is the edge in $G - E$ incident to z . Suppose, on the contrary, that $G - E$ has a Λ -factor, say P . Since $G - \{d, t\}$ is not connected, $G - E - E(D)$ is also not connected. Obviously the component of $G - E - E(D)$ containing z belongs to \mathcal{A} . Therefore by **2.9**, $G - E - E(D)$ has no Λ -factor. Thus P has a 3-vertex path L containing exactly one edge in D adjacent to edge d . Now if C is a component of $G - E - L$, then $v(C) \neq 0 \pmod 3$. Therefore $G - E$ has no Λ -factor, a contradiction.

(p2) Now we prove $(g) \Rightarrow (e)$. Namely, we assume that \dot{E} does not satisfy (e) and we want to show that in this case $G - E$ has a Λ -factor. Let u be the edge of T distinct from a and b .

Suppose that \dot{E} is connected, and so \dot{E} is a 3-edge path. Let V be a 3-vertex path in G containing u and avoiding E . Then $G - V$ has no edges from E , and so $G - V = G - E - V$. By **2.1(a)**, $G - V$ has a Λ -factor.

Now suppose that \dot{E} is not connected, and so \dot{E} has exactly two components induced by $\{a, b\}$ and by c , respectively. Since \dot{E} does not satisfy (e), \dot{E} is not a claw and not a triangle, and so $u, t \notin E$.

(p2.1) Suppose that c belongs to no triangle in G . Since \dot{E} does not satisfy (e), $G - E$ is connected. Clearly, $G - E$ is claw-free. Also $G - E$ has exactly two end-blocks and the block of one edge t is one of them. By **1.10**, $G - E$ has a Λ -factor.

(p2.2) Now suppose that c belongs to a triangle D in G . Then $D \neq T$. Let $G' = G - D - \{a, b\}$. Then G' is claw-free and has no edges from E .

Suppose that G' is connected. Then as in **(p2.1)**, G' has exactly two end-blocks. By **1.10**, G' has a Λ -factor, say P . Let $L = D - c$, and so L is a 3-vertex path. Then

$P \cup \{L\}$ is a Λ -factor in $G - E$.

Now suppose that G' is not connected. Let C be the component of G' containing edge t and q the edge connecting C and D . Let L be a 3-vertex path in G containing q and an edge in $D - c$. It is sufficient to show that $G - E - L$ has a Λ -factor. Obviously $G - E - L$ is claw-free. Let Q be a component of $G - E - L$. Since \dot{E} does not satisfy (e4), $v(Q) = 0 \bmod 3$ and Q has exactly two end-blocks. By **1.10**, Q has a Λ -factor. Therefore $G - E - L$ also has a Λ -factor. \square

From **2.10** we have, in particular:

2.11 *Suppose that $G = F^\Delta$, where F is a cubic 2-connected graph (and so G is claw-free). Let $E \subset E(G)$ and $|E| = 2$. Then $G - E$ has a Λ -factor.*

From **2.10** we also have:

2.12 *Suppose that G is a 3-connected claw-free graph. Let $E \subset E(G)$ and $|E| = 3$. Then $G - E$ has a Λ -factor if and only if \dot{E} is not a claw and not a triangle.*

2.13 *Suppose that G is a 2-connected claw-free graph, $v(G) = 1 \bmod 3$ and $x \in V(G)$. Then $G - x$ has a Λ -factor.*

Proof (uses **1.10**). Let $x \in V(G)$. Since $v(G) = 1 \bmod 3$, clearly $v(G - x) = 0 \bmod 3$. Since G is 2-connected, $G - x$ is connected. Since G is claw-free, $G - x$ is claw-free and has at most two end-blocks. By **1.10**, $G - x$ has a Λ -factor. \square

2.14 *Suppose that G is a 3-connected claw-free graph, $v(G) = 1 \bmod 3$, $x \in V(G)$ and $e \in E(G)$. Then $G - \{x, e\}$ has a Λ -factor.*

Proof (uses **2.8** and **2.13**). Since G is 3-connected, $G - x$ is a 2-connected claw-free graph. Since $v(G) = 1 \bmod 3$, we have $v(G - x) = 0 \bmod 3$. By **2.13**, $G - x$ has a Λ -factor P . If $e \notin E(G - x)$, then P is a Λ -factor of $G - \{x, e\}$. If $e \in E(G - x)$, then by **2.8**, $G - \{x, e\}$ has a Λ -factor. \square

Obviously, the claim in **2.14** may not be true for a claw-free graph of connectivity 2.

From **2.1**, **2.3**, **2.7**, **2.11**, **2.13**, and **2.14** we have, in particular:

2.15 *All claims in **1.9** except for (z5) are true for 3-connected claw-free graphs and (z5) is true for cubic, 2-connected graphs such that every vertex belongs to exactly one triangle.*

References

- [1] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications*, MacMillan Co., New York, 1976.
- [2] R. Deistel, *Graph Theory*, Springer-Verlag, New York, 2005.
- [3] P. Hell and D. Kirkpatrick, Packing by complete bipartite graphs, *SIAM J. Algebraic Discrete Math.* **7** (1986), 199–209.
- [4] A. Kaneko, A. Kelmans, T. Nishimura, On packing 3-vertex paths in a graph, *J. Graph Theory* **36** (2001) 175–197.
- [5] A. Kelmans, Packing k -edge trees in graphs of restricted degrees, *J. Graph Theory* **55** (2007) 306–324 (see also *DIMACS Research Report 2000–44*, Rutgers University (2000)).
- [6] A. Kelmans, Packing a k -path in a cubic graph is NP-hard for $k \geq 3$, manuscript 2000; presented at the DIMACS conference “Graph Partition”, July 2000.
- [7] A. Kelmans, Λ -packings in connected cubic graphs, manuscript 2002; presented at the 34th Southeastern International Conference on Combinatorics, Graph Theory and Computing, March, 2003.
- [8] A. Kelmans, On Λ -packings in 2-connected graphs, *RUTCOR Research Report 21–2005*, Rutgers University (2005).
- [9] A. Kelmans, On Λ -packings in 3-connected graphs, *RUTCOR Research Report 23–2005*, Rutgers University (2005).
- [10] A. Kelmans, Counterexamples to the cubic graph domination conjecture, *ArXiv: math.CO/0607512*, 24 July 2006.
- [11] A. Kelmans and D. Mubayi, How many disjoint 2-edge paths must a cubic graph have ? *Journal of Graph Theory*, **45** (2004) 57–79 (see also *DIMACS Research Report 2000–23*, Rutgers University (2000)).
- [12] D. West, *Introduction to Graph Theory*, Prentice Hall, 2001.